

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6972255>

# Modulation of cytokine expression by traditional medicines: A review of herbal immunomodulators

Article in *Alternative Medicine Review: a Journal of Clinical Therapeutic* · July 2006

Source: PubMed

---

CITATIONS

285

---

READS

3,436

6 authors, including:



[Kevin Spelman](#)

Health, Education & Research in Botanical Medicine

31 PUBLICATIONS 988 CITATIONS

[SEE PROFILE](#)



[Steve Ottersberg](#)

Technica Botanica LLC

1 PUBLICATION 285 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Malaria Research [View project](#)



Method Validation [View project](#)

# Modulation of Cytokine Expression by Traditional Medicines: A Review of Herbal Immunomodulators

Kevin Spelman, MS; JJ Burns, ND; Douglas Nichols, ND; Nasha Winters, ND;  
Steve Ottersberg, MS; Mark Tenborg, ND

## Abstract

**INTRODUCTION:** Modulation of cytokine secretion may offer novel approaches in the treatment of a variety of diseases. One strategy in the modulation of cytokine expression may be through the use of herbal medicines. A class of herbal medicines, known as immunomodulators, alters the activity of immune function through the dynamic regulation of informational molecules such as cytokines. This may offer an explanation of the effects of herbs on the immune system and other tissues. For this informal review, the authors surveyed the primary literature on medicinal plants and their effects on cytokine expression, taking special care to analyze research that utilized the multi-component extracts equivalent to or similar to what are used in traditional medicine, clinical phytotherapy, or in the marketplace. **METHODOLOGY:** MEDLINE, EBSCO, and BIOSIS were used to identify research on botanical medicines, in whole or standardized form, that act on cytokine activity through different models, i.e., *in vivo* (human and animal), *ex vivo*, or *in vitro*. **RESULTS:** Many medicinal plant extracts had effects on at least one cytokine. The most frequently studied cytokines were IL-1, IL-6, TNF, and IFN. *Acalypha wilkesiana*, *Acanthopanax gracilistylus*, *Allium sativum*, *Ananus comosus*, *Cissampelos sympodialis*, *Coriolus versicolor*, *Curcuma longa*, *Echinacea purpurea*, *Grifola frondosa*, *Harpagophytum procumbens*, *Panax ginseng*, *Polygala tenuifolia*, *Poria cocos*, *Silybum marianum*, *Smilax glabra*, *Tinospora cordifolia*, *Uncaria tomentosa*, and *Withania somnifera* demonstrate modulation of multiple

cytokines. **CONCLUSION:** The *in vitro* and *in vivo* research demonstrates that the reviewed botanical medicines modulate the secretion of multiple cytokines. The reported therapeutic success of these plants by traditional cultures and modern clinicians may be partially due to their effects on cytokines. Phytotherapy offers a potential therapeutic modality for the treatment of many differing conditions involving cytokines. Given the activity demonstrated by many of the reviewed herbal medicines and the increasing awareness of the broad-spectrum effects of cytokines on autoimmune conditions and chronic degenerative processes, further study of phytotherapy for cytokine-related diseases and syndromes is warranted. (*Altern Med Rev* 2006;11(2):128-150)

Kevin Spelman, MS, RH(AHG), MCPP – Chair of Clinical Division, Department of Herbal Medicine, Tai Sophia Institute; adjunct research scientist, Department of Chemistry & Biochemistry, University of North Carolina, Greensboro, NC. Correspondence address: Tai Sophia Institute, 7750 Montpelier Road, Laurel, MD 20723  
Email: spelman123@earthlink.net

JJ Burns, ND – 2004 graduate of Southwest College of Naturopathic Medicine, Tempe, AZ; private practice, Scottsdale, AZ.

Mark Tenborg, ND – 2004 graduate of Southwest College of Naturopathic Medicine, Tempe, AZ; environmental health consultant, AZ.

Douglas Nichols, ND – 2004 graduate of Southwest College of Naturopathic Medicine, Tempe, AZ; private practice, Snowflake, AZ.

Nasha Winters, ND, LAc – 2000 graduate of the International Institute of Chinese Medicine in Albuquerque, NM; 2003 graduate of Southwest College of Naturopathic Medicine, Tempe, AZ; private practice, Durango, CO.

Steve Ottersberg, MS Biochemistry – 1999 graduate of Arizona State University; adjunct professor of chemistry at Fort Lewis College, Durango, CO.

## Introduction

Cytokines, a large group of soluble extracellular proteins or glycoproteins, are key intercellular regulators and mobilizers. Classified into family groups (e.g., interleukins, interferons, and chemokines) based on the structural homologies of their receptors, these proteins were initially believed to act primarily as antiviral<sup>1</sup> or antineoplastic<sup>2</sup> agents. They are now seen to be crucial to innate and adaptive inflammatory responses, cell growth and differentiation, cell death, angiogenesis, and developmental as well as repair processes.<sup>3</sup> Their secretion, by virtually every nucleated cell type, is usually an inducible response to injurious stimuli.<sup>3</sup> In addition, cytokines provide a link between organ systems, providing molecular cues for maintaining physiological stability.<sup>4</sup> Medical literature of the last several decades reveals an array of conditions, from cardiovascular disease to frailty, whose onset and course may be influenced by cytokines.<sup>5</sup>

The diverse and far-reaching influences of these proteins can be seen in the central nervous system (CNS); cytokines cause the brain to produce neurochemical, neuroendocrine, neuroimmune, and behavioral shifts.<sup>6</sup> Abnormal cytokine production has been demonstrated in neuropsychiatric disorders such as attention deficit hyperactivity disorder, obsessive-compulsive disorder, and anorexia nervosa.<sup>6,7</sup> Cytokines also appear to play a role in depression, schizophrenia, and Alzheimer's disease,<sup>7</sup> and may be a common link between insomnia and depression.<sup>8,9</sup> In addition, there appears to be an involvement of cytokines in anhedonia (the inability to experience pleasure) and learned helplessness.<sup>10</sup>

The understanding of stimuli that invoke cytokine secretion has expanded. Besides chronic infections, negative emotions and stressful experiences have been shown to stimulate production of proinflammatory cytokines.<sup>5</sup> In addition to involvement in neuropsychiatric disorders, these diverse glycoproteins have activity in all body systems. As models of physiology continue to develop beyond compartmentalized organ systems, elucidation of the global activity of cytokines offers further support to an expanding understanding of cell-to-cell communication. The inflammatory processes of cardiovascular disease are one such example. Beyond leukocytes,

the liver, heart, vessel walls, and adipose tissue are known to produce cytokines; thus any of these tissues may potentially contribute to the inflammatory nature of cardiovascular disease.<sup>11</sup>

As a result of the growing recognition of cytokine activities, altering cytokine expression and targeting their receptors may offer therapeutic potential. Current pharmacological strategies include cytokine antagonist, agonist, inhibition, and stimulation models.<sup>12</sup> Therapeutic application of cytokines in clinical medicine has rapidly surpassed the FDA's 1986 approval of an interferon (IFN) agonist for the treatment of hairy cell leukemia. In 2001, an antagonist to tumor necrosis factor (TNF), a pivotal cytokine in the pathogenesis of rheumatoid arthritis (RA), was described as one of the most important advances in RA treatment.<sup>13</sup> In addition, interleukin-1 $\beta$  (IL-1 $\beta$ ) and TNF antagonists offer options for the treatment of periodontal disease.<sup>14</sup> A novel approach in the treatment of asthma is the inhibition of T-helper 2 (TH2) derived cytokine expression, resulting in downstream effects on IgE and eosinophils.<sup>15</sup> Interleukin-10 (IL-10) demonstrates modulation of brain inflammation, which may have application for conditions such as Alzheimer's disease.<sup>16</sup> In addition, interleukin-2 (IL-2) and interleukin-12 (IL-12) in combination may provide a potential therapeutic approach for neuroblastomas.<sup>17</sup>

Due to their diverse and pleiotropic activities, cytokine treatments may prove promising for disorders seemingly unrelated to immune function. However, much of their therapeutic effect relies on direct influence of immune activity. For example, in the field of oncology, progress has been made in the therapeutic use of several interleukins, including IL-4, -6, -11 and -12.<sup>18</sup> In combination with surgery, pretreatment with IL-2 may enhance survival rates in patients with renal cell carcinoma.<sup>19</sup> IL-18 demonstrates antitumor effects in leukemia.<sup>20</sup> The interferons are used in the treatment of hepatitis B and C, malignant melanoma, follicular lymphoma, and AIDS-related Kaposi's sarcoma.<sup>21</sup>

However, as with the development of many nascent pharmacological strategies, the occurrence of adverse events generates barriers to successful therapeutic applications. Such obstacles have delayed progress in the use of several synthetic cytokines.

Treatment with recombinant cytokines has yielded a number of adverse effects, such as transient lymphopenias induced by IFN, IL-2, and TNF. Monocytopenia has been reported with the use of interferon-gamma (IFN- $\gamma$ ) and TNF, while IL-2, IFN- $\alpha$ , and TNF induce neutrophilia.<sup>22</sup> Patient experience of flu-like symptoms with the use of interferons makes adherence to a therapeutic protocol a challenge. Both IL-2 and IFN- $\alpha$ , used for the treatment of hepatitis C and some cancers, are known to evoke depression, fatigue, sleepiness, irritability, and loss of appetite.<sup>23</sup> These toxic side effects have limited the clinical value of such therapies.<sup>24</sup>

In light of the adverse events experienced with cytokine-targeted therapy, it could prove useful to consider the use of phytotherapy in the modulation of cytokine expression. Immune-related illnesses have long been treated with herbal medicines. The primary literature suggests many of the effects of botanicals may be via cytokine modulation.<sup>25</sup> The term immunomodulator has been used in the phytotherapy literature to describe botanical medicines believed to influence immunity.<sup>26</sup> In regard to phytotherapy, immunomodulators may be defined as botanical medicines that alter the activities of the immune system via the dynamic regulation of informational molecules – cytokines, hormones, neurotransmitters, and other peptides.

This article provides an informal review of the scientific literature regarding the effects of botanical medicines on cytokines. Islam and Carter point out that therapy based on medicinal plants, such as the immunomodulators, is based on diverse constituents or groups of constituents and therefore, researching isolated constituents to reveal modes of activity disregards the principles of phytotherapy.<sup>27</sup> In addition, when clinicians use medicinal plant preparations in practice, they often do not treat with isolated constituents. Therefore, in order to maintain relevance for clinical phytotherapy, this informal survey was limited to herbal medicines available in the marketplace or preparations that represent multi-component botanical medicines.

## Methodology

### Search Strategy

The databases MEDLINE, EBSCO, and BIOSIS were searched for appropriate studies. Titles were screened for all hits to the terms “herbs and cytokines” and “Chinese medicine and cytokines” and “Ayurveda and cytokines.” A language restriction of English was observed.

### Criteria for Inclusion

The following parameters were necessary for study inclusion:

- Investigations on whole herbs (e.g., seed, leaf, root, stem, flower, or entire plant), standardized extracts, or extractions of whole herbs not reduced to one constituent were accepted. Research on isolated constituents or multiple herbal formulations were generally rejected. Fungi, although technically not plants, were included as they are commonly used in phytotherapy.
- All study model types were accepted – *in vitro*, *ex vivo*, and *in vivo* (both animal and human) models were accepted.
- Information on methods of herbal preparation, concentration of the plant preparation, and dose/exposure time were required.
- Only studies demonstrating activity with regard to cytokines were included.

One hundred thirty-nine titles and abstracts were reviewed for inclusion criteria. Ninety-five studies were eliminated due to single constituent-based research or insignificant results. Forty-nine papers met the criteria.

## Results

Information collected as a result of searches is listed in Tables 1-5. The majority of the research used *in vitro* models, but *in vivo* animal models were also utilized. Data in Tables 1A and 1B catalog *in vivo* results, noting the genus and species of the plants, the

Table 1A. *In vivo* Effects of Botanicals on Cytokine Expression

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Aloe secundiflora</i>	Sap	Aqueous	200 mg/kg (pre) 400 mg/kg (post) infection	2 weeks (pre) + 7 days (post)	<i>In vivo</i> , Fowl	IL-6	(Waihenya et al, 2002) <sup>28</sup>
<i>Angelica sylvestris</i>	Root	Aqueous	640 mg/kg PO	2 weeks	<i>In vivo</i> , Murine	TNF	(Haranaka et al, 1985) <sup>29</sup>
<i>Asparagus racemosus</i>	Root	80% Ethanol	100 mg/kg QD PO	17 weeks	<i>In vivo</i> , Murine	IL-1 $\alpha$ TNF	(Dhuley, 1997) <sup>30</sup>
<i>Bupleurum falcatum</i>	Root	Aqueous	640 mg/kg PO	2 weeks	<i>In vivo</i> , Murine	TNF	(Haranaka et al, 1985) <sup>29</sup>
<i>Cinnamomum cassia</i>	Bark	Aqueous	560 mg/kg PO	2 weeks	<i>In vivo</i> , Murine	TNF	(Haranaka et al, 1985) <sup>29</sup>
<i>Cnidium monnieri</i>	Rhizome	Aqueous	560 mg/kg PO	2 weeks	<i>In vivo</i> , Murine	TNF	(Haranaka et al, 1985) <sup>29</sup>
<i>Coptis spp.</i>	Rhizome	1% Coptis pwd in standard mouse diet	1% <i>C. rhizoma</i> in diet	7 days	<i>In vivo</i> , Murine	IL-6	(Iizuka et al, 2000) <sup>31</sup>
<i>Panax ginseng</i>	Root	40% Ethanol	150 mg/kg QD injection	7 days	<i>In vivo</i> , Murine	IL-4	(Song et al, 2002) <sup>32</sup>
<i>Panax ginseng</i>	Root	40% Ethanol	150 mg/kg QD injection	21 days	<i>In vivo</i> , Murine	IL-4 TNF IFN- $\gamma$	(Song et al, 2002) <sup>32</sup>

**Table 1B. *In vivo* Effects of Botanicals on Cytokine Expression**

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Perilla frutescens</i>	Leaf	Aqueous	0.4 mL/murine x2 PO	2 doses 18 hrs+1 hr	<i>In vivo</i> , Murine	TNF	(Ueda and Yamazaki, 1997) <sup>33</sup>
<i>Picrothiza kurroa</i>	Rhizome	80% Ethanol	100 mg/kg QD PO	17 weeks	<i>In vivo</i> , Murine	IL-1 $\alpha$ TNF	(Dhuley, 1997) <sup>30</sup>
<i>Polygala tenuifolia</i>	Root	Aqueous	2x10 <sup>3</sup> mg/kg QD PO	9 days	<i>In vivo</i> , Murine	IL-4 IFN- $\gamma$	(Hong et al, 2002) <sup>34</sup>
<i>Silybum marianum</i>	Seed and Fruit	Silymarin	10 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-2, -4	(Johnson et al, 2003) <sup>35</sup>
<i>Silybum marianum</i>	Seed	Silymarin	250 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-1 $\beta$ , -6 TNF	(Johnson et al, 2003) <sup>35</sup>
<i>Silybum marianum</i>	Seed	Silymarin	250 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-2, -4	(Johnson et al, 2002) <sup>36</sup>
<i>Smilax glabra</i>	Rhizome	Aqueous	400 mg/kg QD PO	14 days	<i>Ex vivo</i> , Murine	IL-1, -2 TNF	(Jiang and Xu, 2003) <sup>37</sup>
<i>Tinospora cordifolia</i>	Root and Herb	80% Ethanol	100 mg/kg QD PO	17 weeks	<i>In vivo</i> , Murine	IL-1 $\alpha$ TNF	(Dhuley, 1997) <sup>30</sup>
<i>Withania somnifera</i>	Root and Leaf	80% Ethanol	100 mg/kg QD PO	17 weeks	<i>In vivo</i> , Murine	IL-1 $\alpha$ TNF	(Dhuley, 1997) <sup>30</sup>
<i>Withania somnifera</i>	Root	70% Methanol	20 mg QD IP	5 days	<i>In vivo</i> , Murine	IL-2 TNF IFN- $\gamma$	(Davis and Kuttan, 1999) <sup>38</sup>
<i>Withania somnifera</i>	Root	70% Methanol	20 mg QD IP	10 days	<i>In vivo</i> , Murine	IL-2 IFN- $\gamma$	(Davis and Kuttan, 1999) <sup>38</sup>

Table 2A. *In vitro* (human cell) Effects of Aqueous Botanical Extracts on Cytokine Expression

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Acanthopanax gracilistylus</i>	Bark	Aqueous	1.08 x 10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Acanthopanax gracilistylus</i>	Bark	Aqueous	20 µg/mL incubation	3 days	<i>In vitro</i> , Human	IL-1, -6 IFN-γ, TNF	(Shan et al, 1999) <sup>40</sup>
<i>Astragalus membranaceus</i>	Root	Aqueous	1.5x10 <sup>5</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-6	(Shon et al, 2002) <sup>41</sup>
<i>Astragalus membranaceus</i>	Root	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Cinnamomum cassia</i>	Bark	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Codonopsis pilosula</i>	Root	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Derris scandens</i>	Stem	Aqueous	10 µg/mL incubation	5 days	<i>In vitro</i> , Human	IL-2	(Sriwanthana and Chavalittumrong, 2001) <sup>42</sup>
<i>Epimedium brevicornum</i>	Herb	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Oldenlandia diffusa</i>	Herb	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Rauwolfia serpentina</i>	Root	Aqueous	5 µg/mL incubation	42 hours	<i>In vitro</i> , Human	TNF	(Jin et al, 2002) <sup>43</sup>
<i>Rauwolfia serpentina</i>	Root	Aqueous	10 µg/mL incubation	42 hours	<i>In vitro</i> , Human	IFN-γ	(Jin et al, 2002) <sup>43</sup>
<i>Schisandra chinensis</i>	Fruit	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>
<i>Sinomenium acutum</i>	Stem	Aqueous	0.1 µg/mL incubation	1 hour	<i>In vitro</i> , Murine	TNF	(Kim et al, 1999) <sup>44</sup>
<i>Smilax glabra</i>	Rhizome	Aqueous	400 mg/kg QD PO	14 days	<i>Ex vivo</i> , Murine	IL-1, -2 TNF	(Jiang and Xu, 2003) <sup>37</sup>
<i>Typhonium sp.</i>	Rhizome	Aqueous	1.08x10 <sup>3</sup> µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1	(Shan et al, 1999) <sup>39</sup>

**Table 2B. *In vitro* (human cell) Effects of Ethanolic Botanical Extracts on Cytokine Expression**

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Curcuma longa</i>	Rhizome	Acetone or Ethanol, % unspecified	1.8 µg/mL incubation	4 hours	<i>In vitro</i> , Human	TNF	(Chan, 1995) <sup>45</sup>
<i>Curcuma longa</i>	Rhizome	Acetone or Ethanol, % unspecified	1.8 µg/mL incubation	18 hours	<i>In vitro</i> , Human	IL-1	(Chan, 1995) <sup>45</sup>
<i>Harpagophytum procumbens</i>	Root	60% Ethanol	>100 µg/mL incubation	0.5 hour	<i>In vitro</i> , Human	IL-1β, -6	(Fiebich et al, 2001) <sup>46</sup>
<i>Harpagophytum procumbens</i>	Root	60% Ethanol	100 µg/mL incubation	0.5 hour	<i>In vitro</i> , Human	TNF	(Fiebich et al, 2001) <sup>46</sup>
<i>Sparassis crispa</i>	Fruit	Defatted by Ethanol, Aqueous	100 µg/mL incubation	48 hours	<i>In vitro</i> , Murine	IL-6	(Harada et al, 2002) <sup>47</sup>
<i>Tripterygium wilfordii</i>	Bark	50% Hot Ethanol	0.6 µg/mL incubation	48 hours	<i>In vitro</i> , Human	IL-2	(Chou and Chang, 1998) <sup>48</sup>
<i>Zingiber officinale</i>	Rhizome	50% Ethanol	2x10 <sup>4</sup> µg/mL incubation	18 hours	<i>In vitro</i> , Human	IL-1β GM-CSF*	(Chang et al, 1995) <sup>49</sup>
<i>Zingiber officinale</i>	Rhizome	50% Ethanol	1x10 <sup>4</sup> µg/mL incubation	18 hours	<i>In vitro</i> , Human	IL-6	(Chang et al, 1995) <sup>49</sup>

\* granulocyte/macrophage-colony stimulating factor



Table 2C. *In vitro* (human cell) Effects of Other Botanical Extracts on Cytokine Expression

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Acalypha wilkesiana</i>	Seed	6.8% Ethanol 5.9% Water 42.5% Hexane	10 µg/mL incubation	2 hours	<i>In vitro</i> , Human	IL-5, -6, IFN-γ, TNF	(Bussing et al, 1999) <sup>50</sup>
<i>Allium sativum</i>	Bulb	Crushed extract	≥0.1 µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-12	(Hodge et al, 2002) <sup>51</sup>
<i>Allium sativum</i>	Bulb	Crushed extract	≥104 µg/mL incubation	24 hours	<i>In vitro</i> , Human	IL-1α, -2, -6, -8, -10, IFN-γ, TNF	(Hodge et al, 2002) <sup>51</sup>
<i>Ampelopsis brevipedunculata</i>	Unspecified	Methanol, then DMSO	100 µg/mL incubation	3 days	<i>In vitro</i> , Human	IL-1β TNF	(Kuo et al, 1999) <sup>52</sup>
<i>Curcuma longa</i>	Rhizome	Curcumin in DMSO 0.1%	18 µg/mL incubation	2 hours	<i>In vitro</i> , Human	IL-8	(Hidaka et al, 2002) <sup>53</sup>
<i>Echinacea purpurea</i>	Flower and Herb	Dried or fresh juice in 20% ethanol	0.012 µg/mL incubation	18 hours	<i>In vitro</i> , Human	IL-1	(Burger et al, 1997) <sup>54</sup>
<i>Echinacea purpurea</i>	Flower and Herb	Dried or fresh juice in 20% ethanol	0.012 µg/mL incubation	72 hours	<i>In vitro</i> , Human	IL-6	(Burger et al, 1997) <sup>54</sup>
<i>Echinacea purpurea</i>	Flower and Herb	Dried or fresh juice in 20% ethanol	0.025 µg/mL incubation	36 hours	<i>In vitro</i> , Human	IL-10	(Burger et al, 1997) <sup>54</sup>
<i>Echinacea purpurea</i>	Flower and Herb	Fresh juice in 20% ethanol	0.012 µg/mL incubation	36 hours	<i>In vitro</i> , Human	TNF	(Burger et al, 1997) <sup>54</sup>
<i>Echinacea purpurea</i>	Flower and Herb	Dried juice in 20% ethanol	0.025 µg/mL incubation	36 hours	<i>In vitro</i> , Human	TNF	(Burger et al, 1997) <sup>54</sup>
<i>Ludwigia octovalvis</i>	Unspecified	Methanol, then DMSO	100 µg/mL incubation	3 days	<i>In vitro</i> , Human	IL-1β TNF	(Kuo et al, 1999) <sup>52</sup>
<i>Rhus semiolata</i>	Unspecified	Methanol, then DMSO	100 µg/mL incubation	48 hours	<i>In vitro</i> , Human	IL-1β TNF	(Kuo et al, 1999) <sup>52</sup>
<i>Tabernaemontana divaricata</i>	Unspecified	Methanol, then DMSO	100 µg/mL incubation	3 days	<i>In vitro</i> , Human	IL-1β TNF	(Kuo et al, 1999) <sup>52</sup>

**Table 3A. *In vitro* (animal cell) Effects of Aqueous Botanical Extracts on Cytokine Expression**

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Acanthopanax senticosus</i>	Seed	Aqueous	10 <sup>3</sup> µg/mL	0.5 hour	<i>In vitro</i> , Murine	TNF	(Yi et al., 2002) <sup>65</sup>
<i>Acer nikoense</i>	Flower	Aqueous	260 µg/mL	1 hour	<i>In vitro</i> , Murine	TNF	(Fujiki et al., 2003) <sup>66</sup>
<i>Cnidium monnieri</i>	Rhizome	Aqueous	560x10 <sup>3</sup> µg/kg PO	2 weeks	<i>In vivo</i> , Murine	TNF	(Haranaka et al., 1985) <sup>29</sup>
<i>Dichroa febrifuga</i>	Root	Aqueous	500 µg/mL incubation	20 hours	<i>In vitro</i> , Murine	TNF	(Kim et al., 2000) <sup>57</sup>
<i>Ixeris dentata</i>	Whole plant	Aqueous	100 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	TNF	(Chung et al., 2002) <sup>68</sup>
<i>Panax quinquefolius</i>	Root	Aqueous	100 µg/mL incubation	20 hours	<i>In vitro</i> , Murine	TNF	(Assineve et al., 2002) <sup>59</sup>
<i>Polygala tenuifolia</i>	Root	Aqueous	1 µg/mL incubation	18 hours	<i>In vitro</i> , Murine	IL-1 TNF	(Kim et al., 1998) <sup>60</sup>
<i>Rosa davurica</i>	Fruit	Aqueous	100 µg/mL incubation	16 hours	<i>In vitro</i> , Murine	TNF	(Kim et al., 1999) <sup>44</sup>
<i>Sinomenium acutum</i>	Stem	Aqueous	0.1 µg/mL incubation	1 hour	<i>In vitro</i> , Murine	TNF	(Kim et al., 2000) <sup>61</sup>
<i>Smilax glabra</i>	Rhizome	Aqueous	400 mg/kg QD PO	14 days	<i>Ex vivo</i> , Murine	IL-1, -2 TNF	(Jiang and Xu, 2003) <sup>37</sup>
<i>Uncaria guianensis</i>	Bark	Aqueous	9.5x10 <sup>-3</sup> µg/mL incubation	19 hours	<i>In vitro</i> , Murine	TNF	(Sandoval et al., 2002) <sup>62</sup>
<i>Uncaria tomentosa</i>	Bark	Aqueous	1x10 <sup>-3</sup> µg/mL incubation	19 hours	<i>In vitro</i> , Murine	TNF	(Sandoval et al., 2002) <sup>62</sup>
<i>Uncaria tomentosa</i>	Bark	Aqueous	100 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IL-1, -6	(Lemaire et al., 1999) <sup>63</sup>
<i>Uncaria tomentosa</i>	Bark	Aqueous	14.1x10 <sup>-3</sup> µg/mL incubation	2 hours	<i>In vitro</i> , Animal	TNF	(Sandoval et al., 2002) <sup>62</sup>

plant parts used, methods of preparation, dose, duration of exposure, model utilized, cytokines affected, and references. Tables 2A-C list the *in vitro* results utilizing human cells, categorized by solvents used for the medicinal plant extractions (A, aqueous; B,

ethanolic; and C, other extractions). Tables 3A-C, similar to Tables 2A-C, list the *in vitro* results utilizing animal cells, categorized by solvents used for the medicinal plant extractions (A, aqueous; B, ethanolic; and C, other extractions). Table 4 illustrates the

research conducted on medicinal mushrooms. Tables 5A-E are categorized by cytokine, matching the cytokine and the direction of effect (upregulation or downregulation) exerted by the particular plant.

A large volume of research was disregarded due to the inclusion criteria. Much of the rejected research was based on isolated constituents. Some research on semi-purified compounds, such as curcumin or bromelain, was included due to their frequent use and availability in commerce.

## Discussion

The majority of the research presented in this review relies on *in vitro* and/or animal models; the authors acknowledge the inadequacies of information derived from such research. Both *in vitro* and animal models may be misleading and often prove to be poor representations of human physiology. The lack of pharmacokinetics in an *in vitro* model brings up questions of the relevance of data gathered from such methodology. In addition, animal models often are misrepresentative of human physiology. Nevertheless, data drawn from such sources, coupled with empirical data from traditional uses of botanical medicines, may provide an insight, however limited, to the mode of activity for many of these herbs. *In vivo* and *in vitro* studies for the listed herbs do suggest that the immunomodulating effects of the botanical medicines reviewed may be due, at least in part, to cytokine modulation. Furthermore, given the broad-spectrum effect of cytokines on cell-to-cell communication, it seems likely some of the other organ systems and tissue effects of these herbal immunomodulators are due to modulation of cytokine expression.

### *Astragalus membranaceus*

The root of *Astragalus membranaceus* is traditionally used in Chinese medicine as a “spleen chi tonic” and for various deficiency and wasting conditions.<sup>79</sup> *A. membranaceus*, in an *in vitro* human model, has been shown to lower IL-6.<sup>41</sup> IL-6 is implicated in a number of inflammatory disorders and as a global marker of impending deterioration.<sup>5</sup> The decrease of IL-6 activity provides a possible rationale for thousands of years of use of this plant in deficiency and wasting diseases. In addition, *Astragalus* is also indicated in shortness of breath and edema,

**Table 3B. *In vitro* (animal cell) Effects of Ethanolic Botanical Extracts on Cytokine Expression**

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Cissampelos sympodialis</i>	Leaf	80% Warm Ethanol	12.5 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IL-4, IFN-γ	(Piuvezam et al, 1999) <sup>64</sup>
<i>Cissampelos sympodialis</i>	Leaf	80% Warm Ethanol	6.25 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IL-10	(Piuvezam et al, 1999) <sup>64</sup>
<i>Embllica officinalis</i>	Fruit	70% Ethanol	250 µg/mL incubation	48 hours	<i>In vitro</i> , Murine	IFN-γ	(Sai Ram et al, 2003) <sup>65</sup>
<i>Sparasssis crispa</i>	Fruit	Defatted by Ethanol Aqueous	100 µg/mL incubation	48 hours	<i>In vitro</i> , Murine	IL-6	(Harada et al, 2002) <sup>47</sup>

**Table 3C. *In vitro* (animal cell) Effects of Other Botanical Extracts on Cytokine Expression**

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Ananas comosus</i>	Stem	Bromelain	50 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	TNF	(Engwerda et al., 2001) <sup>66</sup>
<i>Ananas comosus</i>	Stem	Bromelain	50 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IFN-γ	(Engwerda et al., 2001) <sup>66</sup>
<i>Echinacea purpurea</i>	Unspecified	Simulated digestion	80 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IL-1α, IL-1β	(Rininger et al., 2000) <sup>67</sup>
<i>Echinacea purpurea</i>	Unspecified	Simulated digestion	320 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	IL-6	(Rininger et al., 2000) <sup>67</sup>
<i>Echinacea purpurea</i>	Unspecified	Simulated digestion	5 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	TNF	(Rininger et al., 2000) <sup>67</sup>
<i>Paeonia suffruticosa</i>	Bark	95% Methanol	100 µg/mL incubation	1 hour	<i>In vitro</i> , Bovine	IL-8	(Oh et al., 2003) <sup>68</sup>
<i>Pinus maritima</i>	Bark	Pycnogenol	50 µg/mL incubation	6 hours	<i>In vitro</i> , Murine	IL-1β, -2	(Cho et al., 2001) <sup>69</sup>
<i>Scutellaria baicalensis</i>	Root	70% Methanol	10 µg/mL incubation	24 hours	<i>In vitro</i> , Murine	TNF	(Kim et al., 2001) <sup>70</sup>
<i>Silybum marianum</i>	Seed and Fruit	Methanol, then Hexane	25 or 250 µg/mL	60 hours	<i>In vitro</i> , Murine	IL-4, -10, IFN-γ	(Wilasrusnee et al., 2002) <sup>71</sup>
<i>Silybum marianum</i>	Seed and Fruit	Silymarin	10 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-2, -4	(Johnson et al., 2003) <sup>85</sup>
<i>Silybum marianum</i>	Seed	Silymarin	250 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-1β, -6, TNF	(Johnson et al., 2003) <sup>85</sup>
<i>Silybum marianum</i>	Seed	Silymarin	250 mg/kg QD IP injection	5 days	<i>In vivo</i> , Murine	IL-2, -4	(Johnson et al., 2002) <sup>86</sup>
<i>Terminalia chebula</i>	Fruit	50% Methanol	1.0 mg/mL incubation	6 hours	<i>In vitro</i> , Murine	TNF	(Shin et al., 2001) <sup>72</sup>
<i>Tylophora asthmatica</i>	Leaf	Methanol	19x10 <sup>-6</sup> µg/mL and 19.5x10 <sup>-3</sup> µg/mL incubation	48 hours	<i>In vitro</i> , Murine	IL-2	(Ganguly et al., 2001) <sup>73</sup>
<i>Tylophora asthmatica</i>	Leaf	Methanol	39x10 <sup>-3</sup> µg/mL incubation	18 hours	<i>In vitro</i> , Murine	IL-1	(Ganguly et al., 2001) <sup>73</sup>

symptoms that could be suggestive of cardiovascular effects. Notably, increased levels of IL-6 and C-reactive protein are associated with a significant increase

in cardiovascular-related death.<sup>5,11</sup> Thus, a possible mechanism for the cardiovascular effects of *A. membranaceus* could be due to its reduction of IL-6.

Table 4. The Effect of Medicinal Mushrooms on Cytokine Expression

Genus species	Plant Part	Preparation Used	Dose	Duration of Exposure	Model	Cytokines Affected	Author/Date
<i>Cordyceps cicadae</i>	Fruiting body	50% Methanol, then DMSO	100 µg/mL incubation	3 days	<i>In vitro, Human</i>	IL-2 IFN-γ	(Weng et al, 2002) <sup>74</sup>
<i>Cordyceps cicadae</i>	Larvae	50% Methanol, then DMSO	100 µg/mL incubation	3 days	<i>In vitro, Human</i>	IL-2 IFN-γ	(Weng et al, 2002) <sup>74</sup>
<i>Coriolus versicolor</i>	Mycelia	70% Ethanol	5 µL/mL incubation	3 days	<i>In vitro, Human</i>	IL-1β	(Hsieh et al, 2002) <sup>75</sup>
<i>Coriolus versicolor</i>	Mycelia	70% Ethanol	3 µL/mL incubation	3 days	<i>In vitro, Human</i>	IL-6	(Hsieh et al, 2002) <sup>75</sup>
<i>Coriolus versicolor</i>	Mycelia	Aqueous	5 µL/mL incubation	3 days	<i>In vitro, Human</i>	IL-1β, -8	(Hsieh et al, 2002) <sup>75</sup>
<i>Coriolus versicolor</i>	Mycelia	Aqueous	3 µL/mL incubation	3 days	<i>In vitro, Human</i>	IL-6	(Hsieh et al, 2002) <sup>75</sup>
<i>Ganoderma lucidum</i>	Fruiting body	Aqueous Ethanollic precipitation	12.8 µg/mL incubation	48 hours	<i>In vitro, Murine</i>	IL-12	(Cao and Lin, 2002) <sup>76</sup>
<i>Grifola frondosa</i>	Fruiting body	Ethanol	1x10 <sup>3</sup> µg/mL incubation	18 hours	<i>In vitro, Murine</i>	IL-12	(Kodama et al, 2002) <sup>77</sup>
<i>Grifola frondosa</i>	Fruiting body	Ethanol	5x10 <sup>3</sup> µg/kg body wt	14 days	<i>In vivo, Murine</i>	TNF IFN-γ	(Kodama et al, 2002) <sup>77</sup>
<i>Poria cocos</i>	Sclerotium	50% Hot Ethanol	800 µg/mL incubation	6 or 24 hours	<i>In vitro, Human</i>	IL-1β	(Yu and Tseng, 1996) <sup>78</sup>
<i>Poria cocos</i>	Sclerotium	50% Hot Ethanol	400 µg/mL incubation	6 or 24 hours	<i>In vitro, Human</i>	IL-6	(Yu and Tseng, 1996) <sup>78</sup>
<i>Poria cocos</i>	Sclerotium	50% Hot Ethanol	400 µg/mL incubation	3, 6 or 12 hours	<i>In vitro, Human</i>	TNF	(Yu and Tseng, 1996) <sup>78</sup>
<i>Poria cocos</i>	Sclerotium	50% Hot Ethanol	200 µg/mL incubation	3, 6 or 24 hours	<i>In vitro, Human</i>	TGF-β	(Yu and Tseng, 1996) <sup>78</sup>

Table 5A. Botanical Influences on IL-1

Cytokine	Plant	Model	Direction of Effect	Author/Date
IL-1	<i>Acanthopanax gracilistylus</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Astragalus membranaceus</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Cinnamomum cassia</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Codonopsis pilosula</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Curcuma longa</i>	<i>In vitro</i> , Human	Decrease	(Chan, 1995) <sup>45</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Murine	Increase	(Burger et al, 1997) <sup>54</sup>
	<i>Epimedium brevicornum</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Oldenlandia diffusa</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Polygala tenuifolia</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 1998) <sup>60</sup>
	<i>Schisandra chinensis</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Smilax glabra</i>	<i>Ex vivo</i> , Human	Decrease	(Jiang and Xu, 2003) <sup>37</sup>
	<i>Tylophora asthmatica</i>	<i>In vitro</i> , Murine	Increase	(Ganguly et al, 2001) <sup>73</sup>
	<i>Typhonium sp.</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>39</sup>
	<i>Uncaria tomentosa</i>	<i>In vitro</i> , Murine	Decrease	(Lemaire et al, 1999) <sup>63</sup>
IL-1 $\alpha$	<i>Acanthopanax gracilistylus</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>40</sup>
	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Murine	Increase	(Rininger et al, 2000) <sup>67</sup>
	<i>Picrorhiza kurroa</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
	<i>Tinospora cordifolia</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
	<i>Withania somnifera</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
IL-1 $\beta$	<i>Ampelopsis brevipedunculata</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Coriolus versicolor</i>	<i>In vitro</i> , Human	Increase	(Hsieh et al, 2002) <sup>75</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Murine	Increase	(Rininger et al, 2000) <sup>67</sup>
	<i>Harpagophytum procumbens</i>	<i>In vitro</i> , Human	Decrease	(Fiebich et al, 2001) <sup>46</sup>
	<i>Ludwigia octovalvis</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Pinus maritima</i>	<i>In vitro</i> , Murine	Decrease	(Cho et al, 2001) <sup>69</sup>
	<i>Poria cocos</i>	<i>In vitro</i> , Human	Increase	(Yu and Tseng, 1996) <sup>78</sup>
	<i>Rhus semialata</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Increase	(Johnson et al, 2003) <sup>35</sup>
	<i>Tabernaemontana divaricata</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Zingiber officinale</i>	<i>In vitro</i> , Human	Increase	(Chang et al, 1995) <sup>49</sup>

Table 5B. Botanical Influences on IL-2, -4, and -5

Cytokine	Plant	Model	Direction of Effect	Author/Date
IL-2	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Cordyceps cicadae</i> (Fruit body)	<i>In vitro</i> , Human	Increase	(Weng et al, 2002) <sup>74</sup>
	<i>Cordyceps cicadae</i>	<i>In vitro</i> , Human	Decrease	(Weng et al, 2002) <sup>74</sup>
	<i>Derris scandens</i>	<i>In vitro</i> , Human	Increase	(Sriwanthana and Chavalittumrong, 2001) <sup>42</sup>
	<i>Pinus maritima</i>	<i>In vitro</i> , Murine	Decrease	(Cho et al, 2001) <sup>69</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Decrease	(Johnson et al, 2003) <sup>35</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Decrease	(Johnson et al, 2002) <sup>36</sup>
	<i>Smilax glabra</i>	<i>Ex vivo</i> , Murine	Increase	(Jiang and Xu, 2003) <sup>37</sup>
	<i>Tripterygium wilfordii</i>	<i>In vitro</i> , Human	Decrease	(Chou and Chang, 1998) <sup>48</sup>
	<i>Tylophora asthmatica</i>	<i>In vitro</i> , Murine	Decrease	(Ganguly et al, 2001) <sup>73</sup>
	<i>Tylophora asthmatica</i>	<i>In vitro</i> , Murine	Increase	(Ganguly et al, 2001) <sup>73</sup>
	<i>Withania somnifera</i>	<i>In vivo</i> , Murine	Increase	(Davis and Kuttan, 1999) <sup>38</sup>
IL-4	<i>Cissampelos sympodialis</i>	<i>In vitro</i> , Murine	Increase	(Piuvezam et al, 1999) <sup>64</sup>
	<i>Polygala tenuifolia</i>	<i>In vivo</i> , Murine	Increase	(Hong et al, 2002) <sup>34</sup>
	<i>Panax ginseng</i>	<i>In vivo</i> , Murine	Decrease	(Song et al, 2002) <sup>32</sup>
	<i>Panax ginseng</i>	<i>In vivo</i> , Murine	Increase	(Song et al, 2002) <sup>32</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Decrease	(Johnson et al, 2003) <sup>35</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Decrease	(Johnson et al, 2002) <sup>36</sup>
	<i>Silybum marianum</i>	<i>In vitro</i> , Murine	Increase	(Wilasrusmee et al, 2002) <sup>71</sup>
IL-5	<i>Acalypha wilkesiana</i>	<i>In vitro</i> , Human	Increase	(Bussing et al, 1999) <sup>50</sup>

### *Allium sativum*

*Allium sativum* (garlic), like many of the plants highlighted in this review, demonstrates effects on multiple cytokines. Garlic lowered IL-6 in an *in vitro* human model.<sup>51</sup> Besides the hypocholesterolemic, antioxidant, and ACE-inhibition activity of garlic,<sup>80</sup> the effect on IL-6 may offer further insight into garlic's well-known cardiovascular activity.

In the same model, garlic also lowered the proinflammatory cytokine IL-1. IL-1 has been postulated to be involved in the destruction of pancreatic  $\beta$ -cells<sup>80</sup> and garlic demonstrates hypoglycemic action and amelioration of alloxan-induced diabetes

in murine models.<sup>81</sup> IL-1 inhibition may be partially responsible for this activity.

Because of the potential for garlic to reduce the proinflammatory cytokines IL-1, TNF, and IL-8, and stimulate IL-10 secretion (an antagonist of proinflammatory cytokines), Hodge et al<sup>51</sup> concluded that this effect, along with garlic's antimicrobial activity, may provide potential mechanisms for garlic's use in inflammatory bowel disease.<sup>80</sup> IL-10 demonstrates modulation of the immunopathology of brain inflammatory diseases such as Alzheimer's disease, providing another potential use for garlic as a cytokine modulator.<sup>16</sup>



Table 5C. Botanical Influences on IL-6, -8, -10, and -12

Cytokine	Plant	Model	Direction of Effect	Author/Date
IL-6	<i>Acalypha wilkesiana</i>	<i>In vitro</i> , Human	Increase	(Bussing et al, 1999) <sup>50</sup>
	<i>Acanthopanax gracilistylus</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>40</sup>
	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Aloe secundiflora</i>	<i>In vitro</i> , Murine	Increase	(Waihenya et al, 2002) <sup>28</sup>
	<i>Astragalus membranaceus</i>	<i>In vitro</i> , Human	Decrease	(Shon et al, 2002) <sup>41</sup>
	<i>Coptis spp.</i>	<i>In vivo</i> , Murine	Decrease	(Iizuka et al, 2000) <sup>31</sup>
	<i>Coriolus versicolor</i>	<i>In vitro</i> , Human	Decrease	(Hsieh et al, 2002) <sup>75</sup>
	<i>Coriolus versicolor</i>	<i>In vitro</i> , Human	Increase	(Hsieh et al, 2002) <sup>75</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Human	Increase	(Burger et al, 1997) <sup>54</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Murine	Increase	(Rininger et al, 2000) <sup>67</sup>
	<i>Harpagophytum procumbens</i>	<i>In vitro</i> , Human	Decrease	(Fiebich et al, 2001) <sup>46</sup>
	<i>Poria cocos</i>	<i>In vitro</i> , Human	Increase	(Yu and Tseng, 1996) <sup>78</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Increase	(Johnson et al, 2003) <sup>35</sup>
	<i>Sparassis crispa</i>	<i>In vitro</i> , Murine	Increase	(Harada et al, 2002) <sup>47</sup>
	<i>Uncaria tomentosa</i>	<i>In vitro</i> , Murine	Increase	(Lemaire et al, 1999) <sup>63</sup>
<i>Zingiber officinale</i>	<i>In vitro</i> , Human	Increase	(Chang et al, 1995) <sup>49</sup>	
IL-8	<i>Coriolus versicolor</i>	<i>In vitro</i> , Human	Decrease	(Hsieh et al, 2002) <sup>75</sup>
	<i>Curcuma longa</i> (Curcumin)	<i>In vitro</i> , Human	Decrease	(Hidaka et al, 2002) <sup>53</sup>
	<i>Paeonia suffruticosa</i>	<i>In vitro</i> , Bovine	Decrease	(Oh et al, 2003) <sup>68</sup>
IL-10	<i>Allium sativum</i>	<i>In vitro</i> , Human	Increase	(Hodge et al, 2002) <sup>51</sup>
	<i>Cissampelos sympodialis</i>	<i>In vitro</i> , Murine	Increase	(Piuvezam et al, 1999) <sup>64</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Human	Increase	(Burger et al, 1997) <sup>54</sup>
	<i>Silybum marianum</i>	<i>In vitro</i> , Murine	Increase	(Wilasrusmee et al, 2002) <sup>71</sup>
IL-12	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Ganoderma lucidum</i>	<i>In vitro</i> , Murine	Increase	(Cao and Lin, 2002) <sup>76</sup>
	<i>Grifola frondosa</i>	<i>In vitro</i> , Murine	Increase	(Kodama et al, 2002) <sup>77</sup>

### The Case for Multi-Component Remedies: A Hypothesis

One of the criticisms of botanical medicines is they are “crude drugs” representing a dilute mixture consisting of hundreds of compounds, not concentrated to contain any single active constituent. Laboratory studies clearly elucidate that the overall

pharmacological effects and therapeutic efficacies of medicinal plants often do not derive from a single compound, but from several compounds generating synergic activity.<sup>82-86</sup> A number of researchers have proposed that multi-component pharmacological agents that hit multiple targets impact the complex



Table 5D. Botanical Influences on TNF

Cytokine	Plant	Model	Direction of Effect	Author/Date
TNF	<i>Acalypha wilkesiana</i>	<i>In vitro</i> , Human	Increase	(Bussing et al, 1999) <sup>50</sup>
	<i>Acanthopanax gracilistylus</i>	<i>In vitro</i> , Human	Increase	(Shan et al, 1999) <sup>40</sup>
	<i>Acanthopanax senticosus</i>	<i>In vitro</i> , Murine	Decrease	(Yi et al, 2002) <sup>55</sup>
	<i>Acer nikoense</i>	<i>In vitro</i> , Murine	Decrease	(Fujiki et al, 2003) <sup>56</sup>
	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Ampelopsis brevipedunculata</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Ananus comosus</i> (Bromelain)	<i>In vitro</i> , Murine	Increase	(Engwerda et al, 2001) <sup>66</sup>
	<i>Angelica sylvestris</i>	<i>In vivo</i> , Murine	Increase	(Haranaka et al, 1985) <sup>29</sup>
	<i>Asparagus racemosus</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
	<i>Bupleurum falcatum</i>	<i>In vivo</i> , Murine	Increase	(Haranaka et al, 1985) <sup>29</sup>
	<i>Cinnamomum cassia</i>	<i>In vivo</i> , Murine	Increase	(Haranaka et al, 1985) <sup>29</sup>
	<i>Cnidium monnieri</i>	<i>In vivo</i> , Murine	Increase	(Haranaka et al, 1985) <sup>29</sup>
	<i>Curcuma longa</i>	<i>In vitro</i> , Human	Decrease	(Chan, 1995) <sup>45</sup>
	<i>Dichroa febrifuga</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 2000) <sup>57</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Human	Increase	(Burger et al, 1997) <sup>54</sup>
	<i>Echinacea purpurea</i>	<i>In vitro</i> , Murine	Increase	(Rininger et al, 2000) <sup>67</sup>
	<i>Grifola frondosa</i>	<i>In vivo</i> , Murine	Increase	(Kodama et al, 2002) <sup>77</sup>
	<i>Harpagophytum procumbens</i>	<i>In vitro</i> , Human	Decrease	(Fiebich et al, 2001) <sup>46</sup>
	<i>Ixeris dentate</i>	<i>In vitro</i> , Murine	Increase	(Chung et al, 2002) <sup>58</sup>
	<i>Ludwigia octovalvis</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Panax ginseng</i>	<i>In vivo</i> , Murine	Increase	(Song et al, 2002) <sup>32</sup>
	<i>Panax quinquefolius</i>	<i>In vitro</i> , Murine	Increase	(Assinewe et al, 2002) <sup>59</sup>
	<i>Perilla frutescens</i>	<i>In vivo</i> , Murine	Decrease	(Ueda and Yamazaki, 1997) <sup>33</sup>
	<i>Picrorhiza kurroa</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
	<i>Polygala tenuifolia</i>	<i>In vitro</i> , Murine	Decrease	(Hong et al, 2002) <sup>34</sup>
	<i>Polygala tenuifolia</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 1998) <sup>60</sup>
	<i>Poria cocos</i>	<i>In vitro</i> , Human	Increase	(Yu and Tseng, 1996) <sup>78</sup>
	<i>Rauwolfia serpentina</i>	<i>In vitro</i> , Human	Increase	(Jin et al, 2002) <sup>43</sup>
	<i>Rhus semialata</i>	<i>In vitro</i> , Human	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Rosa davurica</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 1999) <sup>44</sup>
	<i>Scutellaria baicalensis</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 2001) <sup>70</sup>
	<i>Silybum marianum</i>	<i>In vivo</i> , Murine	Increase	(Johnson et al, 2003) <sup>35</sup>
	<i>Silybum marianum</i>	<i>In vitro</i> , Murine	Increase	(Wilasrusmee et al, 2002) <sup>71</sup>
	<i>Sinomenium acutum</i>	<i>In vitro</i> , Murine	Decrease	(Kim et al, 1999) <sup>44</sup>
	<i>Smilax glabra</i>	<i>In vivo</i> , Murine	Decrease	(Jiang and Xu, 2003) <sup>37</sup>
	<i>Tabernaemontana divaricata</i>	<i>In vitro</i> , Murine	Decrease	(Kuo et al, 1999) <sup>52</sup>
	<i>Terminalia chebula</i>	<i>In vitro</i> , Human	Increase	(Shin et al, 2001) <sup>72</sup>
	<i>Tinospora cordifolia</i>	<i>In vivo</i> , Murine	Decrease	(Dhuley, 1997) <sup>30</sup>
	<i>Uncaria tomentosa</i>	<i>In vitro</i> , Murine	Decrease	(Sandoval et al, 2002) <sup>62</sup>
	<i>Withania somnifera</i>	<i>In vivo</i> , Murine	Increase	(Dhuley, 1997) <sup>30</sup>
<i>Withania somnifera</i>	<i>In vivo</i> , Murine	Decrease	(Davis and Kuttan, 1999) <sup>38</sup>	

Table 5E. Botanical Influences on TGF- $\beta$ , IFN- $\gamma$ , and GM-CSF

Cytokine	Plant	Model	Direction of Effect	Author/Date
TGF- $\beta$	<i>Poria cocos</i>	<i>In vitro</i> , Human	Decrease	(Yu and Tseng, 1996) <sup>78</sup>
IFN- $\gamma$	<i>Acalypha wilkesiana</i>	<i>In vitro</i> , Human	Increase	(Bussing et al, 1999) <sup>50</sup>
	<i>Acanthopanax gracilistylus</i>	<i>In vitro</i> , Human	Decrease	(Shan et al, 1999) <sup>40</sup>
	<i>Allium sativum</i>	<i>In vitro</i> , Human	Decrease	(Hodge et al, 2002) <sup>51</sup>
	<i>Ananus comosus</i> (Bromelain)	<i>In vitro</i> , Murine	Increase	(Engwerda et al, 2001) <sup>66</sup>
	<i>Cissampelos sympodialis</i>	<i>In vitro</i> , Murine	Decrease	(Piuvezam et al, 1999) <sup>64</sup>
	<i>Cordyceps cicada</i> (Fruit Body)	<i>In vitro</i> , Human	Increase	(Weng et al, 2002) <sup>74</sup>
	<i>Cordyceps cicada</i> (Larvae)	<i>In vitro</i> , Human	Decrease	(Weng et al, 2002) <sup>74</sup>
	<i>Embllica officinalis</i>	<i>In vitro</i> , Murine	Increase	(Sai Ram et al, 2003) <sup>65</sup>
	<i>Grifola frondosa</i>	<i>In vivo</i> , Murine	Increase	(Kodama et al, 2002) <sup>77</sup>
	<i>Panax ginseng</i>	<i>In vivo</i> , Murine	Increase	(Song et al, 2002) <sup>32</sup>
	<i>Polygala tenuifolia</i>	<i>In vivo</i> , Murine	Decrease	(Hong et al, 2002) <sup>34</sup>
<i>Rauwolfia serpentina</i>	<i>In vitro</i> , Murine	Increase	(Jin et al, 2002) <sup>43</sup>	
<i>Withania somnifera</i>	<i>In vivo</i> , Murine	Increase	(Davis and Kuttan, 1999) <sup>38</sup>	
GM-CSF*	<i>Zingiber officinalis</i>	<i>In vitro</i> , Human	Increase	(Chang, 1995) <sup>49</sup>

\* granulocyte/macrophage-colony stimulating factor

equilibrium of whole cellular networks more favorably than drugs that act on a single target.<sup>87-93</sup> Keith and Zimmerman<sup>91</sup> suggest many genes might need complementary action to modify disease processes. In other words, efficacious therapy might depend on perturbing more than one target. In addition, multi-target agents need affect their targets only partially, which corresponds well with the presumed low-affinity, substrate/enzyme interactions of medicinal plants.<sup>87-91</sup> The partial “perturbations” of medicinal plants on a pharmacological network may accurately mimic physiological scenarios where hundreds of different enzyme systems and receptor types and subtypes are triggered simultaneously.<sup>87</sup> This is compared to the complete elimination of a single network node (enzyme or receptor system), which is a rather unusual phenomenon not typically found in a physiological scenario.<sup>87</sup> Clinicians have historically overcome such single target insufficiency by using combination drug therapy; for example, therapeutic application of drug cocktails are increasingly utilized in AIDS, cancer, and resistant infections.

Substantial historical, empirical, and scientific evidence demonstrates that whole plants, not just isolated constituents, have immunomodulating activity. Combinations of phytochemicals and cytokines may also provide a novel approach to clinical medicine. Engwerda et al<sup>66</sup> demonstrated the potential for combination therapy using bromelain, a mixture of cysteine proteases from the stems of pineapple plants. In this model, bromelain alone showed limited activity on cytokine secretions. However, if combined with cytokines, a synergic effect was observed. Bromelain with IFN- $\gamma$  significantly enhanced TNF production beyond the effect of IFN- $\gamma$  alone. In addition, when bromelain was combined with IL-12, a significant increase of IFN- $\gamma$  was demonstrated compared to that of only IL-12.<sup>66</sup> Since such responses could enhance acquired immune responses in addition to innate immune responses, critical for first-line defense against many infectious agents, such combinations are likely important.<sup>94</sup> For example, combination therapies could act as vaccine adjuvants, enhancing their efficacy.<sup>95-97</sup>

The combination of medicinal plants with one another or other pharmacological agents fits well into a phytotherapeutic paradigm. Commonly, many of these constituents have additive or synergic activity, while a class of constituents or a single constituent may potentiate a single pharmacologically active molecule.<sup>83,98</sup>

Csermely<sup>89</sup> suggests that a pharmacological strategy directed toward multiple targets could result in more efficient therapeutic outcomes. Broader specificity, lower affinity, multi-component compounds, as found in botanical medicines, can be more efficient than high affinity, high specificity compounds.<sup>87</sup> Moreover, the use of whole plants, instead of isolated chemicals, may offer a safer clinical strategy in the treatment of many diseases.<sup>85,99</sup> Network models of pharmacology, which view human physiology as a complex web of molecular interactions, strongly imply that herbal remedies serve clinical therapy efficaciously, efficiently, and safely.

Equally, this web-like nature is reflected in the immune system by the concerted signaling of cytokines. Cytokines operate both as a cascade and as a network, regulating the production of other cytokines and cytokine receptors, while stimulating the production of acute-phase proteins.<sup>100</sup> Endogenous levels of cytokines are in the nanomolar to picomolar range, suggesting that dilute mixtures of biologically active compounds may provide therapeutic benefit. Illustrating the therapeutic potential for dilute mixtures of biologically active compounds, a group of researchers found subclinical doses of oral IFN- $\alpha$  can provide powerful, broad-spectrum benefits.<sup>1</sup>

In another study, when cytokine levels were compared to symptoms in individuals with cardiovascular disease, Testa et al<sup>101</sup> demonstrated that circulating levels of cytokines increased with severity of symptoms. Considering the variety of adverse events listed for recombinant cytokine therapies, perhaps subtle perturbations of the cytokine network should be considered. The dilute nature of botanical immunomodulators may offer a reasonable strategy for subtle induction of a variety of cytokines.

Most likely, cells are seldom exposed to only a single cytokine. Rather, combinations of cytokines and other messenger molecules generate biologically relevant informational cues.<sup>100</sup> This is demonstrated

by the synergic antitumor effects observed from combining IL-12 gene therapy with other cytokines, chemokines, or co-stimulatory molecules.<sup>24</sup> The effects of cytokines on their target cells and tissues may be inhibited or enhanced by other cytokines, hormones, and cytokine-receptor antagonists and circulating receptors. Just as pharmacological activity by specific plant constituents is suggested to be affected by combinations of constituents,<sup>83,98,102</sup> combinations of cytokines have been found to have additive, inhibitory, or synergic effects.<sup>100</sup> Further research may find that the herbal immunomodulators affecting multiple cytokines can each generate a unique signature of immune perturbation dependent on the concerted effect on arrays of cytokines.

### *Biphasic Effects*

Both exogenous and endogenous compounds can have opposing, dose-dependent biological effects. For example, Calabrese and Baldwin discuss biphasic aortic smooth muscle response to an adrenergic agonist; low doses of isoproterenol bind  $\beta$ -adrenergic receptors, inducing relaxation of aortic smooth muscle. However, at higher doses, where the  $\beta$ -receptors are saturated and the  $\alpha$ -receptors are also bound, isoproterenol induces aortic constriction.<sup>103</sup> Similarly, Sapolsky<sup>104</sup> discusses the effects of glucocorticoids (GCs) on performance of hippocampal-dependent memory, suggesting low-to-moderate levels of endogenous GCs, saturating mineralocorticoid receptors (and some GC receptors), could enhance this process, while higher doses of GCs impair memory.

This biphasic effect can be noted in Table 5D. For example, *Withania somnifera* (ashwagandha) has been found to influence the expression of TNF. Dhuley et al<sup>30</sup> found *W. somnifera* increased TNF expression, while Davis and Khutan<sup>38</sup> showed it decreased TNF.

The models used in these two laboratory investigations are disparate. Dhuley<sup>30</sup> used the carcinogen ochratoxin A (OTA) against murine macrophages to suppress chemotactic activity induced by IL-1 and TNF. Ashwagandha, at an oral dose of 100 mg/kg daily, countered the immunosuppressive effects of OTA, raising TNF expression and theoretically restoring chemotactic activity.

In contrast, Davis and Khutan,<sup>38</sup> using a dose of 20 mg/animal daily by IP injection, found TNF was lowered in the *W. somnifera* group without an inducer. Although the two models are unrelated by dose, duration of exposure, and method of administration, the question still arises as to the paradoxical effects on TNF secretion. The inconsistency in the TNF results could lie in the utilization of divergent models, although these authors suggest the possibility of a biphasic dose response.

TNF is believed to be a key factor in cancer anorexia-cachexia syndrome.<sup>105</sup> Ashwagandha has a history of thousands of years of use in the treatment of wasting syndromes and general debility,<sup>106</sup> and is often currently used clinically as an adjuvant to cancer treatments.<sup>26,107</sup>

Known as anthrapachaka in the Ayurvedic system, *Tylophora asthmatica* is traditionally used in the treatment of asthma, allergies, and autoimmune disorders.<sup>108</sup> *Tylophora* demonstrates a biphasic effect on IL-2 secretion. Ganguly et al demonstrated this effect in an *in vitro* model. Using the same model throughout their investigations, a lower dose of *T. asthmatica* increased IL-2 levels, while more than a thousand-fold increase in dose reduced IL-2 levels, demonstrating a paradoxical response to the same exogenous stimulus.<sup>73</sup>

## Conclusion

Although many of the plants listed in this review appear to affect only a few cytokines, it is the lead author's opinion that future research will further demonstrate the broad-spectrum activity of herbal medicine. Currently, the research on the influence of botanical medicines on cytokines and other messenger molecules is limited. Informational molecules and many of their receptors may likely turn out to be modulated by plants, both herbal medicines and foods, providing potential for future therapeutics.

Despite the fact that the majority of research in this review was performed with *in vitro* or animal models, there is substantial historical, empirical, and scientific evidence that whole plants, not just isolated constituents, have immunomodulating activity. The *in vitro* and *in vivo* research suggests that the reviewed botanical medicines modulate cytokines, and that such modulation may provide the mechanism of

action for many of their therapeutic effects. Further research (particularly clinical studies) is indicated to elucidate the effects of botanical medicines and to support or refute the hypotheses presented in this article.

## References

1. Young AS, Cummins JM. The history of interferon and its use in animal therapy. *East Afr Med J* 1990;67:SS31-SS63.
2. Terlikowski SJ. Tumour necrosis factor and cancer treatment: a historical review and perspectives. *Rocz Akad Med Bialymst* 2001;46:5-18.
3. Oppenheim JJ. Cytokines: past, present, and future. *Int J Hematol* 2001;74:3-8.
4. O'Sullivan RL, Lipper G, Lerner EA. The neuro-immuno-cutaneous-endocrine network: relationship of mind and skin. *Arch Dermatol* 1998;134:1431-1435.
5. Kiecolt-Glaser JK, McGuire L, Robles TF, Glaser R. Psychoneuroimmunology and psychosomatic medicine: back to the future. *Psychosom Med* 2002;64:15-28.
6. Kronfol Z, Remick DG. Cytokines and the brain: implications for clinical psychiatry. *Am J Psychiatry* 2000;157:683-694.
7. Mittleman BB. Cytokine networks in Sydenham's chorea and PANDAS. *Adv Exp Med Biol* 1997;418:933-935.
8. Dubas-Slemp H, Marmurowska-Michalowska H, Szuster-Ciesielska A, et al. The role of cytokines in depression. *Psychiatr Pol* 2003;37:787-798. [Article in Polish]
9. Lustberg L, Reynolds CF. Depression and insomnia: questions of cause and effect. *Sleep Med Rev* 2000;4:253-262.
10. Capuron L, Dantzer R. Cytokines and depression: the need for a new paradigm. *Brain Behav Immun* 2003;17:S119-S124.
11. Rader DJ. Inflammatory markers of coronary risk. *N Engl J Med* 2000;343:1179-1182.
12. Sommer C. Animal studies on neuropathic pain: the role of cytokines and cytokine receptors in pathogenesis and therapy. *Schmerz* 1999;13:315-323.
13. Keystone EC. Tumor necrosis factor-alpha blockade in the treatment of rheumatoid arthritis. *Rheum Dis Clin North Am* 2001;27:427-443.
14. Takashiba S, Naruishi K, Murayama Y. Perspective of cytokine regulation for periodontal treatment: fibroblast biology. *J Periodontol* 2003;74:103-110.
15. Stirling RG, Chung KF. Future treatments of allergic diseases and asthma. *Br Med Bull* 2000;56:1037-1053.



16. Szczepanik AM, Ringheim GE. IL-10 and glucocorticoids inhibit Abeta(1-42)- and lipopolysaccharide-induced pro-inflammatory cytokine and chemokine induction in the central nervous system. *J Alzheimers Dis* 2003;5:105-117.
17. Siapati KE, Barker S, Kinnon C, et al. Improved antitumor immunity in murine neuroblastoma using a combination of IL-2 and IL-12. *Br J Cancer* 2003;88:1641-1648.
18. Oleksowicz L, Dutcher JP. A review of the new cytokines: IL-4, IL-6, IL-11, and IL-12. *Am J Ther* 1994;1:107-115.
19. Bohm M, Ittenson A, Klatte T, et al. Pretreatment with interleukin-2 modulates perioperative immunodysfunction in patients with renal cell carcinoma. *Folia Biol (Praha)* 2003;49:63-68.
20. Akamatsu S, Arai N, Hanaya T, et al. Antitumor activity of interleukin-18 against the murine T-cell leukemia/lymphoma EL-4 in syngeneic mice. *J Immunother* 2002;25:S28-S34.
21. Waldmann TA. Immunotherapy: past, present and future. *Nat Med* 2003;9:269-277.
22. Engel A, Kern WV, Murdter G, Kern P. Kinetics and correlation with body temperature of circulating interleukin-6, interleukin-8, tumor necrosis factor alpha and interleukin-1 beta in patients with fever and neutropenia. *Infection* 1994;22:160-164.
23. Wichers M, Maes M. The psychoneuroimmunopathophysiology of cytokine-induced depression in humans. *Int J Neuropsychopharmacol* 2002;5:375-388.
24. Belardelli F, Ferrantini M. Cytokines as a link between innate and adaptive antitumor immunity. *Trends Immunol* 2002;23:201-208.
25. Spelman K. Immunomodulators of the Ayurvedic Pharmacopoeia. Paper presented at: Southwest Conference on Botanical Medicine; April 6-7, 2002; Southwest College of Naturopathic Medicine & Health Sciences; Tempe, AZ.
26. Spelman K. Immunomodulatory Botanicals as Adjuvant Therapy in Cancer: Case Studies. Paper presented at: Southwest Conference on Botanical Medicine; April 8-9, 2000; Southwest College of Naturopathic Medicine & Health Sciences; Tempe, AZ.
27. Islam J, Carter R. Use of Echinacea in upper respiratory tract infection. *South Med J* 2005;98:311-318.
28. Waihenya RK, Mtambo MM, Nkwengulila G, Minga UM. Efficacy of crude extract of *Aloe secundiflora* against *Salmonella gallinarum* in experimentally infected free-range chickens in Tanzania. *J Ethnopharmacol* 2002;79:317-323.
29. Haranaka K, Satomi N, Sakurai A, et al. Antitumor activities and tumor necrosis factor producibility of traditional Chinese medicines and crude drugs. *Cancer Immunol Immunother* 1985;20:1-5.
30. Dhuley JN. Effect of some Indian herbs on macrophage functions in ochratoxin A treated mice. *J Ethnopharmacol* 1997;58:15-20.
31. Iizuka N, Miyamoto K, Hazama S, et al. Anticachectic effects of *Coptidis rhizoma*, an anti-inflammatory herb, on esophageal cancer cells that produce interleukin 6. *Cancer Lett* 2000;158:35-41.
32. Song Z, Wu H, Mathee K, et al. Gerimax ginseng regulates both humoral and cellular immunity during chronic *Pseudomonas aeruginosa* lung infection. *J Altern Complement Med* 2002;8:459-466.
33. Ueda H, Yamazaki M. Inhibition of tumor necrosis factor-alpha production by orally administering a perilla leaf extract. *Biosci Biotechnol Biochem* 1997;61:1292-1295.
34. Hong T, Jin GB, Yoshino G, et al. Protective effects of Polygalae root in experimental TNBS-induced colitis in mice. *J Ethnopharmacol* 2002;79:341-346.
35. Johnson VJ, He Q, Osuchowski MF, Sharma RP. Physiological responses of a natural antioxidant flavonoid mixture, silymarin, in BALB/c mice: III. Silymarin inhibits T-lymphocyte function at low doses but stimulates inflammatory processes at high doses. *Planta Med* 2003;69:44-49.
36. Johnson VJ, Osuchowski MF, He Q, Sharma RP. Physiological responses to a natural antioxidant flavonoid mixture, silymarin, in BALB/c mice: II. Alterations in thymic differentiation correlate with changes in c-myc gene expression. *Planta Med* 2002;68:961-965.
37. Jiang J, Xu Q. Immunomodulatory activity of the aqueous extract from rhizome of *Smilax glabra* in the later phase of adjuvant-induced arthritis in rats. *J Ethnopharmacol* 2003;85:53-59.
38. Davis L, Kuttan G. Effect of *Withania somnifera* on cytokine production in normal and cyclophosphamide treated mice. *Immunopharmacol Immunotoxicol* 1999;21:695-703.
39. Shan BE, Yoshida Y, Sugiura T, Yamashita U. Stimulating activity of Chinese medicinal herbs on human lymphocytes *in vitro*. *Int J Immunopharmacol* 1999;21:149-159.
40. Shan BE, Yoshida Y, Sugiura T, Yamashita U. Suppressive effect of Chinese medicinal herb, *Acanthopanax gracilistylus*, extract on human lymphocytes *in vitro*. *Clin Exp Immunol* 1999;118:41-48.

41. Shon YH, Kim JH, Nam KS. Effect of *Astragali radix* extract on lipopolysaccharide-induced inflammation in human amnion. *Biol Pharm Bull* 2002;25:77-80.
42. Sriwanthana B, Chavalittumrong P. *In vitro* effect of *Derris scandens* on normal lymphocyte proliferation and its activities on natural killer cells in normals and HIV-1 infected patients. *J Ethnopharmacol* 2001;76:125-129.
43. Jin GB, Hong T, Inoue S, et al. Augmentation of immune cell activity against tumor cells by *Rauwolfia radix*. *J Ethnopharmacol* 2002;81:365-372.
44. Kim HM, Park YA, Lee EJ, Shin TY. Inhibition of immediate-type allergic reaction by *Rosa davurica* Pall. in a murine model. *J Ethnopharmacol* 1999;67:53-60.
45. Chan MM. Inhibition of tumor necrosis factor by curcumin, a phytochemical. *Biochem Pharmacol* 1995;49:1551-1556.
46. Fiebich BL, Heinrich M, Hiller KO, Kammerer N. Inhibition of TNF-alpha synthesis in LPS-stimulated primary human monocytes by Harpagophytum extract SteiHap 69. *Phytomedicine* 2001;8:28-30.
47. Harada T, Miura N, Adachi Y, et al. Effect of SCG, 1,3-beta-D-glucan from *Sparassis crispa* on the hematopoietic response in cyclophosphamide induced leukopenic mice. *Biol Pharm Bull* 2002;25:931-939.
48. Chou CT, Chang SC. The inhibitory effect of common traditional anti-rheumatic herb formulas on prostaglandin E and interleukin 2 *in vitro*: a comparative study with *Tripterygium wilfordii*. *J Ethnopharmacol* 1998;62:167-171.
49. Chang CP, Chang JY, Wang FY, Chang JG. The effect of Chinese medicinal herb *Zingiberis rhizoma* extract on cytokine secretion by human peripheral blood mononuclear cells. *J Ethnopharmacol* 1995;48:13-19.
50. Bussing A, Stein GM, Herterich-Akinpelu I, Pfuller U. Apoptosis-associated generation of reactive oxygen intermediates and release of pro-inflammatory cytokines in human lymphocytes and granulocytes by extracts from the seeds of *Acalypha wilkesiana*. *J Ethnopharmacol* 1999;66:301-309.
51. Hodge G, Hodge S, Han P. *Allium sativum* (garlic) suppresses leukocyte inflammatory cytokine production *in vitro*: potential therapeutic use in the treatment of inflammatory bowel disease. *Cytometry* 2002;48:209-215.
52. Kuo YC, Sun CM, Tsai WJ, et al. Blocking of cell proliferation, cytokines production and genes expression following administration of Chinese herbs in the human mesangial cells. *Life Sci* 1999;64:2089-2099.
53. Hidaka H, Ishiko T, Furuhashi T, et al. Curcumin inhibits interleukin 8 production and enhances interleukin 8 receptor expression on the cell surface: impact on human pancreatic carcinoma cell growth by autocrine regulation. *Cancer* 2002;95:1206-1214.
54. Burger RA, Torres AR, Warren RP, et al. Echinacea-induced cytokine production by human macrophages. *Int J Immunopharmacol* 1997;19:371-379.
55. Yi JM, Hong SH, Kim JH, et al. Effect of *Acanthopanax senticosus* stem on mast cell-dependent anaphylaxis. *J Ethnopharmacol* 2002;79:347-352.
56. Fujiki H, Suganuma M, Kurusu M, et al. New TNF-alpha releasing inhibitors as cancer preventive agents from traditional herbal medicine and combination cancer prevention study with EGCG and sulindac or tamoxifen. *Mutat Res* 2003;523-524:119-125.
57. Kim YH, Ko WS, Ha MS, et al. The production of nitric oxide and TNF-alpha in peritoneal macrophages is inhibited by *Dichroa febrifuga* Lour. *J Ethnopharmacol* 2000;69:35-43.
58. Chung HS, Jeong HJ, Han MJ, et al. Nitric oxide and tumor necrosis factor-alpha production by *Ixeris dentata* in mouse peritoneal macrophages. *J Ethnopharmacol* 2002;82:217-222.
59. Assinewe VA, Amason JT, Aubry A, et al. Extractable polysaccharides of *Panax quinquefolius* L. (North American ginseng) root stimulate TNFalpha production by alveolar macrophages. *Phytomedicine* 2002;9:398-404.
60. Kim HM, Lee EH, Na HJ, et al. Effect of *Polygala tenuifolia* root extract on the tumor necrosis factor-alpha secretion from mouse astrocytes. *J Ethnopharmacol* 1998;61:201-208.
61. Kim HM, Moon PD, Chae HJ, et al. The stem of *Sinomenium acutum* inhibits mast cell-mediated anaphylactic reactions and tumor necrosis factor-alpha production from rat peritoneal mast cells. *J Ethnopharmacol* 2000;70:135-141.
62. Sandoval M, Okuhama NN, Zhang XJ, et al. Anti-inflammatory and antioxidant activities of cat's claw (*Uncaria tomentosa* and *Uncaria guianensis*) are independent of their alkaloid content. *Phytomedicine* 2002;9:325-337.

63. Lemaire I, Assinewe V, Cano P, et al. Stimulation of interleukin-1 and -6 production in alveolar macrophages by the neotropical liana, *Uncaria tomentosa* (una de gato). *J Ethnopharmacol* 1999;64:109-115.
64. Piuvezam MR, Pecanha LM, Alexander J, Thomas G. *Cissampelos sympodioides* Eichl. leaf extract increases the production of IL-10 by concanavalin-A-treated BALB/c spleen cells. *J Ethnopharmacol* 1999;67:93-101.
65. Sai Ram M, Neetu D, Deepti P, et al. Cytoprotective activity of Amla (*Emblica officinalis*) against chromium (VI) induced oxidative injury in murine macrophages. *Phytother Res* 2003;17:430-433.
66. Engwerda CR, Andrew D, Murphy M, Mynott TL. Bromelain activates murine macrophages and natural killer cells *in vitro*. *Cell Immunol* 2001;210:5-10.
67. Rininger JA, Kickner S, Chigurupati P, et al. Immunopharmacological activity of Echinacea preparations following simulated digestion on murine macrophages and human peripheral blood mononuclear cells. *J Leukoc Biol* 2000;68:503-510.
68. Oh GS, Pae HO, Choi BM, et al. Inhibitory effects of the root cortex of *Paeonia suffruticosa* on interleukin-8 and macrophage chemoattractant protein-1 secretions in U937 cells. *J Ethnopharmacol* 2003;84:85-89.
69. Cho KJ, Yun CH, Packer L, Chung AS. Inhibition mechanisms of bioflavonoids extracted from the bark of *Pinus maritima* on the expression of proinflammatory cytokines. *Ann N Y Acad Sci* 2001;928:141-156.
70. Kim YO, Leem K, Park J, et al. Cytoprotective effect of *Scutellaria baicalensis* in CA1 hippocampal neurons of rats after global cerebral ischemia. *J Ethnopharmacol* 2001;77:183-188.
71. Wilasrusmee C, Kittur S, Shah G, et al. Immunostimulatory effect of *Silybum marianum* (milk thistle) extract. *Med Sci Monit* 2002;8:BR439-BR443.
72. Shin TY, Jeong HJ, Kim DK, et al. Inhibitory action of water soluble fraction of *Terminalia chebula* on systemic and local anaphylaxis. *J Ethnopharmacol* 2001;74:133-140.
73. Ganguly T, Badheka LP, Sainis KB. Immunomodulatory effect of *Tylophora indica* on Con A induced lymphoproliferation. *Phytomedicine* 2001;8:431-437.
74. Weng SC, Chou CJ, Lin LC, et al. Immunomodulatory functions of extracts from the Chinese medicinal fungus *Cordyceps cicadae*. *J Ethnopharmacol* 2002;83:79-85.
75. Hsieh TC, Kunicki J, Darzynkiewicz Z, Wu JM. Effects of extracts of *Coriolus versicolor* (I'm-Yunity) on cell-cycle progression and expression of interleukins-1 beta, -6, and -8 in promyelocytic HL-60 leukemic cells and mitogenically stimulated and nonstimulated human lymphocytes. *J Altern Complement Med* 2002;8:591-602.
76. Cao LZ, Lin ZB. Regulation on maturation and function of dendritic cells by *Ganoderma lucidum* polysaccharides. *Immunol Lett* 2002;83:163-169.
77. Kodama N, Komuta K, Sakai N, Nanba H. Effects of D-Fraction, a polysaccharide from *Grifola frondosa* on tumor growth involve activation of NK cells. *Biol Pharm Bull* 2002;25:1647-1650.
78. Yu SJ, Tseng J. Fu-Ling, a Chinese herbal drug, modulates cytokine secretion by human peripheral blood monocytes. *Int J Immunopharmacol* 1996;18:37-44.
79. Bensky D, Clavey S, Steoger E. *Chinese Herbal Medicine: Materia Medica*. 3rd ed. Seattle, WA: Eastland Press; 2004.
80. Ebadi MS. *Pharmacodynamic Basis of Herbal Medicine*. Boca Raton, FL: CRC Press; 2002.
81. Huang KC, Williams WM. *The Pharmacology of Chinese Herbs*. 2nd ed. Boca Raton: CRC Press; 1999.
82. Spelman K, Duke JA, Bogenschutz-Godwin MJ. The synergy principle in plants, pathogens, insects, herbivores and humans. In: Kaufman PB, ed. *Natural Products from Plants*. Boca Raton, FL: CRC Press; 2006:486
83. Spelman K, Duke JA, Bogenschutz-Godwin MJ. The synergy principle in plants, pathogens, insects, herbivores and humans. In: Kaufman PB, ed. *Natural Products from Plants*. Boca Raton, FL: CRC Press; 2006:478.
84. Wagner H. Phytomedicine research in Germany. *Environ Health Perspect* 1999;107:779-781.
85. Williamson EM. Synergy and other interactions in phytomedicines. *Phytomedicine* 2001;8:401-409.
86. Spinella M. The importance of pharmacological synergy in psychoactive herbal medicines. *Altern Med Rev* 2002;7:130-137.
87. Agoston V, Csermely P, Pongor S. Multiple weak hits confuse complex systems: a transcriptional regulatory network as an example. *Phys Rev E Stat Nonlin Soft Matter Phys* 2005;71:1-8.
88. Csermely P. Strong links are important, but weak links stabilize them. *Trends Biochem Sci* 2004;29:331-334.
89. Csermely P, Agoston V, Pongor S. The efficiency of multi-target drugs: the network approach might help drug design. *Trends Pharmacol Sci* 2005;26:178-182.

90. Keith CT, Borisyy AA, Stockwell BR. Multicomponent therapeutics for networked systems. *Nat Rev Drug Discov* 2005;4:71-78.
91. Keith CT, Zimmermann GR. Multi-target lead discovery for networked systems. *Curr Drug Discov* 2004;19-23.
92. Briskin DP. Medicinal plants and phytomedicines. Linking plant biochemistry and physiology to human health. *Plant Physiol* 2000;124:507-514.
93. Werner E. In silico multicellular systems biology and minimal genomes. *Drug Discov Today* 2003;8:1121-1127.
94. Valiante NM, O'Hagan DT, Ulmer JB. Innate immunity and biodefence vaccines. *Cell Microbiol* 2003;5:755-760.
95. Boyaka PN, Tafaro A, Fischer R, et al. Therapeutic manipulation of the immune system: enhancement of innate and adaptive mucosal immunity. *Curr Pharm Des* 2003;9:1965-1972.
96. Robson NC, Beacock-Sharp H, Donachie AM, Mowat AM. The role of antigen-presenting cells and interleukin-12 in the priming of antigen-specific CD4+ T cells by immune stimulating complexes. *Immunology* 2003;110:95-104.
97. Villinger F. Cytokines as clinical adjuvants: how far are we? *Expert Rev Vaccines* 2003;2:317-326.
98. Gilbert B, Alves LF. Synergy in plant medicines. *Curr Med Chem* 2003;10:13-20.
99. Ernst E. Herbal medicines put into context. *BMJ* 2003;327:881-882.
100. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 1999;340:448-454.
101. Testa M, Yeh M, Lee P, et al. Circulating levels of cytokines and their endogenous modulators in patients with mild to severe congestive heart failure due to coronary artery disease or hypertension. *J Am Coll Cardiol* 1996;28:964-971.
102. Wills RB, Bone K, Morgan M. Herbal products: active constituents, modes of action and quality control. *Nutr Res Rev* 2000;13:47-77.
103. Calabrese EJ, Baldwin LA. Defining hormesis. *Hum Exp Toxicol* 2002;21:91-97.
104. Sapolsky RM. Stress and plasticity in the limbic system. *Neurochem Res* 2003;28:1735-1742.
105. Lelli G, Montanari M, Gilli G, et al. Treatment of the cancer anorexia-cachexia syndrome: a critical reappraisal. *J Chemother* 2003;15:220-225.
106. Upton R, Booth SJ. *Ashwagandha Root, Withania somnifera: Analytical, Quality Control, and Therapeutic Monograph*. Scotts Valley, CA: American Herbal Pharmacopoeia; 2000.
107. Yance DR, Valentine A. *Herbal Medicine, Healing & Cancer*. Los Angeles, CA: Keats Publishing; 1999.
108. Bone K. *Clinical Applications of Ayurvedic and Chinese Herbs, Monographs for the Western Herbal Practitioner*. Warwick, Australia: Phytotherapy Press; 1996.